

HIGHER ORDER MIXING IN PARABOLIC DYNAMICS

HENK BRUIN

The focus of ergodic theory is the study of measure-preserving dynamical systems which, despite their deterministic nature, display chaotic, “random-like” behaviours. One of the fundamental manifestations of chaos in measure-preserving dynamical systems is *mixing*¹. In probabilistic language, mixing means that any two observables become asymptotically independent under the action of the dynamics, or, in other words, their correlations decay. More in general, a measure-preserving system is *k-mixing* if the multiple correlations between any *k* observables decay. Although it has been proven that several mixing systems are also mixing of all orders (i.e., *k-mixing* for all $k \geq 2$), the general question whether 2-mixing implies mixing of all orders remains a famous open problem in ergodic theory (the so-called *Rohlin Problem*).

Once that a given system is known to be mixing or *k-mixing*, it is of fundamental importance, especially for applications, to estimate *how fast* the correlations between smooth observables decay. This has been done successfully for several “highly chaotic” (*hyperbolic*) systems and for some *parabolic* ones. The adjective *parabolic* refers to those dynamical systems which display an intermediate regime between ordered ergodic (elliptic) systems and highly chaotic (hyperbolic) ones. Important examples of such systems are smooth area-preserving flows on surfaces, unipotent flows on quotients of semisimple Lie groups, nilflows on nilmanifolds, and their smooth perturbations, in particular their time-changes. Parabolic flows are known to be typically mixing, but with a slow (namely, polynomial) rate, see, e.g., [1, 5, 6, 7, 8, 10, 11, 16, 17, 19, 20] and references therein for a complete panorama.

The aim of the present proposal for a PhD research project is to advance the current knowledge on mixing properties, in particular on higher order mixing, for smooth parabolic flows, both homogeneous and non-homogeneous. There have been several new developments in this direction in recent years, but the current state of the art is far from satisfactory. We believe that the open questions we describe in this proposal can be addressed by a PhD student and positive answers will attract attention from the scientific community working in parabolic dynamics.

Specific projects:

- (1) The *horocycle flow* is arguably the prime example of a homogeneous parabolic flow. Sharp mixing estimates were established by Ratner in her seminal paper [16]. Mixing of all orders was proved by Marcus [14], and a polynomial upper bound for *k-mixing*, for all $k \geq 3$, follows from a result of Björklund, Einsiedler and Gorodnik [2], which holds in great generality. However, in this specific setting, one could aim at obtaining sharper estimates for higher order mixing. A possible approach to tackle this problem is to combine Ratner’s strategy in [16] with either [2] or [14].
- (2) The exponent governing the rate of decay of *k-correlations* in [2] is not explicit. It would be interesting to consider specific cases, e.g., for unipotent flows on quotients M of $SL_n(\mathbb{R})$, and estimate it for given $k \geq 3$ (in terms of the spectral gap of M). Other flows with slow mixing (stemming from billiards or mixtures of billiards and geodesic flows) exist as well, see e.g. [3, 4], but also here the rates of *k-mixing* for $k \geq 3$ are unknown.
- (3) The study of non-homogeneous parabolic flows has received considerable attention in the last few years. Notably, researchers have tried to generalise the results known for the unipotent flows to their smooth perturbations, in particular to their time-changes (see, among others, [1, 9, 10, 12, 15]). Compared to the unperturbed setting, much less is known on their quantitative mixing properties. Even for time-changes of horocycle flows, optimal rates of mixing are not known. Possible objectives include improving the known estimates on mixing rates for time-changes of unipotent flows, obtaining lower bounds, and studying the case of more general perturbations beyond the case of time-changes (see, e.g., [18]).

REFERENCES

- [1] A. Avila, G. Forni, D. Ravotti and C. Ulcigrai, *Mixing for smooth time-changes of general nilflows*, Adv. Math., **385** (2021), 107759.

¹Sometimes referred to as *strong mixing* or *strong 2-mixing*.

- [2] M. Björklund, M. Einsiedler and A. Gorodnik, *Quantitative multiple mixing*, J. Eur. Math. Soc., **22** (2020), 1475–1529.
- [3] H. Bruin, *On Sinai billiards on flat surfaces with non-flat horns*, Journ. Stat. Phys. **183** (2021).
- [4] H. Bruin, *On volume preserving almost Anosov flows*, Monatsh. Math. appeared electronically on 20/12/2022.
- [5] J. Chaika and A. Wright, *A smooth mixing flow on a surface with nondegenerate fixed points*, J. Amer. Math. Soc., **32** (2019), 81–117.
- [6] B. Fayad, *Polynomial decay of correlations for a class of smooth flows on the two torus*, Bull. Soc. Math. France, **129** (2001), 487–503.
- [7] B. Fayad, *Analytic mixing reparametrizations of irrational flows*, Ergod. Theory Dyn. Syst., **22** (2002), 437–468.
- [8] B. Fayad, G. Forni and A. Kanigowski, *Lebesgue spectrum of countable multiplicity for conservative flows on the torus*, J. Amer. Math. Soc., **34** (2021), 747–813.
- [9] G. Forni and A. Kanigowski, *Multiple mixing and disjointness for time changes of bounded-type Heisenberg nilflows*, J. Éc. Polytech. Math., **7** (2020), 63–91.
- [10] G. Forni and C. Ulcigrai, *Time-changes of horocycle flows*, J. Mod. Dyn., **6** (2012), 251–273.
- [11] A. Kanigowski, J. Kulaga-Przymus and C. Ulcigrai, *Multiple mixing and parabolic divergence in smooth area-preserving flows on higher genus surfaces*, J. Eur. Math. Soc., **21** (2019), 3797–3855.
- [12] A. Kanigowski, M. Lemańczyk and C. Ulcigrai, *On disjointness properties of some parabolic flows*, Invent. Math., **221** (2020), 1–111.
- [13] A. Kanigowski and D. Ravotti, *Polynomial 3-mixing for smooth time-changes of horocycle flows*, Discrete Contin. Dyn. Syst., **40** (2020), 5347–5371.
- [14] B. Marcus, *The horocycle flow is mixing of all degrees*, Invent. Math., **46** (1978), 201–209.
- [15] M. Ratner, *Rigidity of time changes for horocycle flows*, Acta Math., **156** (1986), 1–32. 7:267–288, 1987.
- [16] M. Ratner, *The rate of mixing for geodesic and horocycle flows*, Ergodic Theory Dynam. Systems **7** (1987), 267–288.
- [17] D. Ravotti, *Quantitative mixing for locally Hamiltonian flows with saddle loops on compact surfaces*, Annales Henri Poincaré, **18** (2017), 3815–3861.
- [18] D. Ravotti, *Parabolic perturbations of unipotent flows on compact quotients of $SL(3, \mathbb{R})$* , Commun. Math. Phys., **371** (2019), 331–351.
- [19] D. Ravotti, *Polynomial mixing for time-changes of unipotent flows*, Ann. Sc. norm. super. Pisa Cl. sci., **23** (2022), 1491–1506 .
- [20] C. Ulcigrai, *Mixing of asymmetric logarithmic suspension flows over interval exchange transformations*, Ergodic Theory Dynam. Systems, **27** (2007), 991–1035.